

# Asymptotically Efficient Adaptive Allocation Rules\*

T. L. LAI AND HERBERT ROBBINS

*Department of Statistics, Columbia University, New York, New York 10027*

## 1. INTRODUCTION

Let  $\Pi_j$  ( $j = 1, \dots, k$ ) denote statistical populations (treatments, manufacturing processes, etc.) specified respectively by univariate density functions  $f(x; \theta_j)$  with respect to some measure  $\nu$ , where  $f(\cdot; \cdot)$  is known and the  $\theta_j$  are unknown parameters belonging to some set  $\Theta$ . Assume that  $\int_{-\infty}^{\infty} |x| f(x; \theta) d\nu(x) < \infty$  for all  $\theta \in \Theta$ . How should we sample  $x_1, x_2, \dots$  sequentially from the  $k$  populations in order to achieve the greatest possible expected value of the sum  $S_n = x_1 + \dots + x_n$  as  $n \rightarrow \infty$ ? Starting with [3] there has been a considerable literature on this subject, which is often called the multi-armed bandit problem. The name derives from an imagined slot machine with  $k \geq 2$  arms. (Ordinary slot machines with one arm are one-armed bandits, since in the long run they are as effective as human bandits in separating the victim from his money.) When an arm is pulled, the player wins a random reward. For each arm  $j$  there is an unknown probability distribution  $\Pi_j$  of the reward. The player wants to choose at each stage one of the  $k$  arms, the choice depending in some way on the record of previous trials, so as to maximize the long-run total expected reward. A more worthy setting for this problem is in the context of sequential clinical trials, where there are  $k$  treatments of unknown efficacy to be used in treating a long sequence of patients.

An *adaptive allocation rule*  $\varphi$  is a sequence of random variables  $\varphi_1, \varphi_2, \dots$  taking values in the set  $\{1, \dots, k\}$  and such that the event  $\{\varphi_n = j\}$  ("sample from  $\Pi_j$  at stage  $n$ ") belongs to the  $\sigma$ -field  $\mathcal{F}_{n-1}$  generated by the previous values  $\varphi_1, x_1, \dots, \varphi_{n-1}, x_{n-1}$ . Let  $\mu(\theta) = \int_{-\infty}^{\infty} xf(x; \theta) d\nu(x)$ .

\*Research supported by the National Science Foundation and the National Institutes of Health. This paper was delivered at the Statistical Research Conference at Cornell University, July 6-9, 1983, in memory of Jack Kiefer and Jacob Wolfowitz.

Then

$$\begin{aligned} ES_n &= \sum_{i=1}^n \sum_{j=1}^k E\left(E\left[x_i I_{\{\varphi_i=j\}} \mid \mathcal{F}_{i-1}\right]\right) \\ &= \sum_{j=1}^k \mu(\theta_j) ET_n(j), \end{aligned} \quad (1.1)$$

where

$$T_n(j) = \sum_{i=1}^n I_{\{\varphi_i=j\}} \quad (1.2)$$

is the number of times that  $\varphi$  samples from  $\Pi_j$  up to stage  $n$ . The problem of maximizing  $ES_n$  is therefore equivalent to that of minimizing the "regret"

$$R_n(\theta_1, \dots, \theta_k) = n\mu^* - ES_n = \sum_{j: \mu(\theta_j) < \mu^*} (\mu^* - \mu(\theta_j)) ET_n(j), \quad (1.3)$$

where by definition

$$\mu^* = \max\{\mu(\theta_1), \dots, \mu(\theta_k)\} = \mu(\theta^*) \quad \text{for some } \theta^* \in \{\theta_1, \dots, \theta_k\}. \quad (1.4)$$

Let  $I(\theta, \lambda)$  denote the Kullback-Leibler number

$$I(\theta, \lambda) = \int_{-\infty}^{\infty} [\log(f(x; \theta)/f(x; \lambda))] f(x; \theta) d\nu(x). \quad (1.5)$$

Then  $0 \leq I(\theta, \lambda) \leq \infty$ , and we shall always assume that  $f(\cdot; \cdot)$  is such that

$$0 < I(\theta, \lambda) < \infty \quad \text{whenever } \mu(\lambda) > \mu(\theta), \quad (1.6)$$

and

$$\begin{aligned} &\forall \epsilon > 0 \text{ and } \forall \theta, \lambda \text{ such that } \mu(\lambda) > \mu(\theta), \quad \exists \delta = \delta(\epsilon, \theta, \lambda) > 0 \\ &\text{for which } |I(\theta, \lambda) - I(\theta, \lambda')| < \epsilon \text{ whenever } \mu(\lambda) \leq \mu(\lambda') \leq \mu(\lambda) + \delta. \end{aligned} \quad (1.7)$$

In Sections 3 and 4 we construct adaptive allocation rules  $\varphi$  such that for any fixed values  $\theta_1, \dots, \theta_k$  for which the  $\mu(\theta_j)$  are not all equal,

$$R_n(\theta_1, \dots, \theta_k) \sim \left\{ \sum_{j: \mu(\theta_j) < \mu^*} (\mu^* - \mu(\theta_j)) / I(\theta_j, \theta^*) \right\} \log n \quad \text{as } n \rightarrow \infty. \quad (1.8)$$

(Here and in the sequel we use the notations  $\mu^*$  and  $\theta^*$  defined in (1.4).) Note that by (1.7),  $I(\theta_j, \theta^*) = I(\theta_j, \lambda)$  whenever  $\mu(\theta_j) < \mu(\theta^*) = \mu(\lambda)$ .

The asymptotic behavior (1.8) of the regret will be shown in Sect. 2 to be *optimal* in the sense of

**THEOREM 1.** *Assume that  $I(\theta, \lambda)$  satisfies (1.6) and (1.7), and that  $\Theta$  is such that*

$$\forall \lambda \in \Theta \text{ and } \forall \delta > 0, \exists \lambda' \in \Theta \text{ such that } \mu(\lambda) < \mu(\lambda') < \mu(\lambda) + \delta. \quad (1.9)$$

Let  $\varphi$  be a rule whose regret satisfies, for each fixed  $\theta = (\theta_1, \dots, \theta_k)$ , the condition that as  $n \rightarrow \infty$

$$R_n(\theta) = o(n^a) \quad \text{for every } a > 0. \quad (1.10)$$

Then for every  $\theta$  such that the  $\mu(\theta_j)$  are not all equal,

$$\liminf_{n \rightarrow \infty} R_n(\theta) / \log n \geq \sum_{j: \mu(\theta_j) < \mu^*} (\mu^* - \mu(\theta_j)) / I(\theta_j, \theta^*). \quad (1.11)$$

Condition (1.10) of Theorem 1 implies that for every  $\theta$

$$\lim_{n \rightarrow \infty} n^{-1} ES_n = \mu^*. \quad (1.12)$$

We shall call rules that satisfy (1.12) *consistent*. Under the assumptions of Theorem 1, we shall call rules that satisfy (1.8) whenever the  $\mu(\theta_j)$  are not all equal *asymptotically efficient*. In the case  $k = 2$  Robbins [3] proposed a simple procedure for constructing consistent rules. Let  $a_1 = 1 < a_2 < \dots$  and  $b_1 = 2 < b_2 < \dots$  be any two disjoint, increasing sequences of positive integers such that  $a_n/n \rightarrow \infty$  and  $b_n/n \rightarrow \infty$  as  $n \rightarrow \infty$ . At stage  $n$ , sample from  $\Pi_1$  if  $n \in \{a_1, a_2, \dots\}$ , sample from  $\Pi_2$  if  $n \in \{b_1, b_2, \dots\}$ , and if  $n \notin \{a_1, a_2, \dots, b_1, b_2, \dots\}$  sample from  $\Pi_1$  or  $\Pi_2$  according as the arithmetic mean of all previous observations from  $\Pi_1$  exceeds or does not exceed the arithmetic mean of all previous observations from  $\Pi_2$ . The consistency of this rule follows easily from the strong law of large numbers.

The sequences  $a_n$  and  $b_n$  above are assumed to be prescribed in advance, and a natural question is how to choose them so that  $n^{-1}ES_n$  approaches  $\mu^*$  as rapidly as possible. However, the choice clearly involves the unknown parameters  $\theta_1, \dots, \theta_k$ . It is therefore desirable to let the sequences  $a_n, b_n$  be generated adaptively from the data rather than prescribed in advance. Such an approach was recently followed by Reimnitz [2] who, in the case of two Bernoulli populations, constructed an allocation rule with regret  $R_n(\theta_1, \theta_2) = O(\log n)$ . His rule, however, does not attain the asymptotically optimal rate (1.8).

In this paper we develop a new approach for constructing rather simple rules that are asymptotically efficient. Our approach is based on a certain class of upper confidence bounds, and the idea is described in general in Section 3. Applications to the special cases of normal, Bernoulli, Poisson, and exponential populations are discussed in Section 4.

## 2. A LOWER BOUND FOR THE EXPECTED SAMPLE SIZE FROM AN INFERIOR POPULATION

Let  $\theta = (\theta_1, \dots, \theta_k)$  and let  $P_\theta$  denote the probability measure under which  $\theta_j$  is the parameter corresponding to population  $\Pi_j$ ,  $j = 1, \dots, k$ . Define for  $j = 1, \dots, k$  the parameter sets

$$\begin{aligned}\Theta_j &= \left\{ \theta: \mu(\theta_j) < \max_{i \neq j} \mu(\theta_i) \right\} && \text{("}\theta_j \text{ is not best")}, \\ \Theta_j^* &= \left\{ \theta: \mu(\theta_j) > \max_{i \neq j} \mu(\theta_i) \right\} && \text{("}\theta_j \text{ is the unique best")}. \end{aligned} \quad (2.1)$$

The main result of this section is given by

**THEOREM 2.** *Assume that  $I(\theta, \lambda)$  satisfies (1.6) and (1.7) and that  $\Theta$  satisfies (1.9). Fix  $j \in \{1, \dots, k\}$ , and define  $\Theta_j$  and  $\Theta_j^*$  by (2.1). Let  $\varphi$  be any rule such that for every  $\theta \in \Theta_j^*$ , as  $n \rightarrow \infty$*

$$\sum_{i \neq j} E_\theta T_n(i) = o(n^a) \quad \text{for every } a > 0, \quad (2.2)$$

where  $T_n(i)$ , defined in (1.2), is the number of times that the rule  $\varphi$  samples from  $\Pi_i$  up to stage  $n$ . Then for every  $\theta \in \Theta_j$  and every  $\epsilon > 0$ ,

$$\lim_{n \rightarrow \infty} P_\theta \left\{ T_n(j) \geq (1 - \epsilon)(\log n) / I(\theta_j, \theta^*) \right\} = 1, \quad (2.3)$$

where  $\theta^*$  is defined in (1.4), and hence

$$\liminf_{n \rightarrow \infty} E_\theta T_n(j) / \log n \geq 1 / I(\theta_j, \theta^*).$$

*Proof.* To fix the ideas let  $j = 1$ ,  $\theta \in \Theta_1$ , and  $\theta^* = \theta_2$ . Then  $\mu(\theta_2) > \mu(\theta_1)$  and  $\mu(\theta_2) \geq \mu(\theta_i)$  for  $3 \leq i \leq k$ . Fix any  $0 < \delta < 1$ . In view of (1.6), (1.7), and (1.9), we can choose  $\lambda \in \Theta$  such that

$$\mu(\lambda) > \mu(\theta_2) \quad \text{and} \quad |I(\theta_1, \lambda) - I(\theta_1, \theta_2)| < \delta I(\theta_1, \theta_2). \quad (2.4)$$

Define the new parameter vector  $\gamma = (\lambda, \theta_2, \dots, \theta_k)$ . Then  $\gamma \in \Theta_1^*$ , so by (2.2)

$$E_\gamma(n - T_n(1)) = \sum_{h \neq 1} E_\gamma(T_n(h)) = o(n^a)$$

with  $0 < a < \delta$ , and therefore

$$\begin{aligned} (n - O(\log n)) P_\gamma \{ T_n(1) < (1 - \delta)(\log n)/I(\theta_1, \lambda) \} \\ \leq E_\gamma(n - T_n(1)) = o(n^a). \end{aligned}$$

Letting  $Y_1, Y_2, \dots$  denote successive observations from  $\Pi_1$ , and defining  $L_m = \sum_{i=1}^m \log(f(Y_i; \theta_1)/f(Y_i; \lambda))$ , it follows that

$$\begin{aligned} P_\gamma(C_n) &= o(n^{a-1}), \quad \text{where} \\ C_n &= \{ T_n(1) < (1 - \delta)(\log n)/I(\theta_1, \lambda) \text{ and } L_{T_n(1)} \leq (1 - a)\log n \}. \end{aligned} \quad (2.5)$$

Note that

$$\begin{aligned} P_\gamma \{ T_n(1) = n_1, \dots, T_n(k) = n_k, L_{n_1} \leq (1 - a)\log n \} \\ = \int_{\{T_n(1)=n_1, \dots, T_n(k)=n_k, L_{n_1} \leq (1-a)\log n\}} \prod_{i=1}^{n_1} \frac{f(Y_i; \lambda)}{f(Y_i; \theta_1)} dP_\theta \\ \geq \exp(-(1 - a)\log n) \\ \cdot P_\theta \{ T_n(1) = n_1, \dots, T_n(k) = n_k, L_{n_1} \leq (1 - a)\log n \}. \end{aligned} \quad (2.6)$$

Since  $C_n$  is a disjoint union of events of the form  $\{ T_n(1) = n_1, \dots, T_n(k) = n_k, L_{n_1} \leq (1 - a)\log n \}$  with  $n_1 + \dots + n_k = n$  and  $n_1 < (1 - \delta)(\log n)/I(\theta_1, \lambda)$ , it follows from (2.5) and (2.6) that as  $n \rightarrow \infty$

$$P_\theta(C_n) \leq n^{1-a} P_\gamma(C_n) \rightarrow 0. \quad (2.7)$$

By the strong law of large numbers,  $L_m/m \rightarrow I(\theta_1, \lambda) > 0$ , and therefore  $\max_{i \leq m} L_i/m \rightarrow I(\theta_1, \lambda)$ , a.s.  $[P_\theta]$ . Since  $1 - a > 1 - \delta$ , it then follows that

$$P_\theta \{ L_i > (1 - a)\log n \text{ for some } i < (1 - \delta)(\log n)/I(\theta_1, \lambda) \} \rightarrow 0 \\ \text{as } n \rightarrow \infty. \quad (2.8)$$

From (2.7) and (2.8) we see that

$$\lim_{n \rightarrow \infty} P_\theta \{ T_n(1) < (1 - \delta)(\log n)/I(\theta_1, \lambda) \} = 0.$$

In view of (2.4), this implies that

$$\lim_{n \rightarrow \infty} P_\theta \{ T_n(1) < (1 - \delta)(\log n)/[(1 + \delta)I(\theta_1, \theta_2)] \} = 0,$$

from which (2.3) for  $j = 1$  follows.

*Proof of Theorem 1.* In view of condition (1.10) on the rule  $\varphi$ , it follows from Theorem 2 that for any fixed  $\theta = (\theta_1, \dots, \theta_k)$ , if  $\mu(\theta_j) < \mu^*$  then

$$\liminf_{n \rightarrow \infty} \{ E_{\theta} T_n(j) / \log n \} \geq 1 / I(\theta_j, \theta^*). \quad (2.9)$$

Since  $R_n(\theta) = \sum_{j: \mu(\theta_j) < \mu^*} (\mu^* - \mu(\theta_j)) E_{\theta} T_n(j)$ , (1.11) follows from (2.9).

### 3. CONSTRUCTION OF ASYMPTOTICALLY EFFICIENT ALLOCATION RULES

We describe here a general method of constructing adaptive allocation rules that attain the asymptotic lower bound for the regret given by the right-hand side of (1.11). We first outline briefly the motivation of our approach. In order to attain asymptotic efficiency we shall sample from the population with the largest estimated mean, provided that we have sampled enough from each population to be reasonably confident that this population is indeed superior. Our degree of confidence will depend on the number  $n$  of observations that we have taken so far, and as  $n$  increases we should be increasingly confident that we are not sampling from an inferior population. In view of Theorem 2, if  $\varphi$  is an asymptotically efficient rule, then the number of observations that  $\varphi$  takes from any inferior population  $\Pi_j$  up to stage  $n$  is about  $(\log n) / I(\theta_j, \theta^*)$ . Thus, at stage  $n$ , we need about  $(\log n) / I(\theta_j, \theta^*)$  observations from  $\Pi_j$  to be reasonably confident that it is not a contender. These considerations suggest the following modified "sample-from-the-leader" rule. First, define the "leader" at stage  $n$  as the population with the largest estimated mean among all populations that have been sampled at least  $\delta n$  times, for some predetermined positive number  $\delta < 1/k$ . While we would like to sample from this apparently superior population, we need to make sure that the other populations have been sampled enough for us to be reasonably confident that they are indeed inferior. One way of doing this is to compare certain upper confidence bounds for the mean of an apparently inferior population with the estimated mean of the leader. These confidence bounds are required to satisfy conditions (3.1), (3.2), and (3.3) below to ensure that the allocation rule obtained thereby is asymptotically efficient.

To fix the ideas, let  $Y_1, Y_2, \dots$  be i.i.d. random variables with a common density function  $f(y; \theta)$  with respect to some measure  $\nu$ , where  $\theta \in \Theta$  denotes an unknown parameter. We shall use "upper confidence bounds" for the mean  $\mu(\theta)$ , defined by Borel functions  $g_{ni}: R^i \rightarrow R$  ( $n = 1, 2, \dots$ ;

$i = 1, \dots, n)$  such that for every  $\theta \in \Theta$ ,

$$P_\theta \{ r \leq g_{ni}(Y_1, \dots, Y_i) \text{ for all } i \leq n \} = 1 - o(n^{-1}) \quad \text{for every } r < \mu(\theta), \quad (3.1)$$

$$\lim_{\epsilon \downarrow 0} \left( \limsup_{n \rightarrow \infty} \sum_{i=1}^n P_\theta \{ g_{ni}(Y_1, \dots, Y_i) \geq \mu(\lambda) - \epsilon \} / \log n \right) \leq 1/I(\theta, \lambda) \quad \text{whenever } \mu(\lambda) > \mu(\theta), \quad (3.2)$$

and

$$g_{ni} \text{ is nondecreasing in } n \geq i \text{ for every fixed } i = 1, 2, \dots \quad (3.3)$$

Examples of such confidence bounds in the special cases of normal, Bernoulli, exponential, and Poisson distributions are given in Section 4, along with a general method for their construction.

In addition to these sequences of upper confidence bounds for  $\mu(\theta)$ , we shall also use point estimates  $h_i(Y_1, \dots, Y_i)$  of  $\mu(\theta)$ , where the  $h_i: R^i \rightarrow R$  ( $i = 1, 2, \dots$ ) are Borel functions such that

$$h_i \leq g_{ni} \quad \text{for all } n \geq i, \quad (3.4)$$

and for every  $\theta \in \Theta$ ,

$$P_\theta \left\{ \max_{\delta n \leq i \leq n} |h_i(Y_1, \dots, Y_i) - \mu(\theta)| > \epsilon \right\} = o(n^{-1}) \quad \text{for all } \epsilon > 0 \text{ and } 0 < \delta < 1. \quad (3.5)$$

Note that (3.5) holds for the sample mean  $h_i(Y_1, \dots, Y_i) = (Y_1 + \dots + Y_i)/i$  under the assumption that  $E_\theta Y_1^2 < \infty$  (cf. the proof of Theorem 1 of [1]).

We now make use of the functions  $g_{ni}$  and  $h_i$  to construct an asymptotically efficient rule for sampling  $x_1, x_2, \dots$  sequentially from populations  $\Pi_1, \dots, \Pi_k$  with respective density functions  $f(x; \theta_1), \dots, f(x; \theta_k)$ . For  $j = 1, \dots, k$ , let  $T_n(j)$  denote the number of times that the rule samples from  $\Pi_j$  up to stage  $n$ , and let  $Y_{j1}, \dots, Y_{j, T_n(j)}$  denote the successive observations from  $\Pi_j$  up to stage  $n$ . Define

$$\begin{aligned} \hat{\mu}_n(j) &= h_{T_n(j)}(Y_{j1}, \dots, Y_{j, T_n(j)}), \\ U_n(j) &= g_{n, T_n(j)}(Y_{j1}, \dots, Y_{j, T_n(j)}), \end{aligned} \quad (3.6)$$

and let  $0 < \delta < 1/k$ . To begin with, at stage  $j = 1, 2, \dots, k$ , the rule takes one observation from  $\Pi_j$ . Now suppose that the rule has taken  $n \geq k$

observations. Since  $T_n(1) + \dots + T_n(k) = n$ , we can choose  $j_n \in \{1, \dots, k\}$  such that

$$\hat{\mu}_n(j_n) = \max\{\hat{\mu}_n(j) : T_n(j) \geq \delta n\}. \quad (3.7)$$

At stage  $n + 1$ , writing  $n + 1 = km + j$ , where  $m$  is a positive integer and  $j \in \{1, \dots, k\}$ , we take an observation from  $\Pi_j$  only if

$$\hat{\mu}_n(j_n) \leq U_n(j), \quad (3.8)$$

and sample from  $\Pi_j$  otherwise. This sampling rule will be denoted by  $\varphi^*$ .

The population  $\Pi_{j_n}$  defined by (3.7) can be regarded as the "leader" at the end of stage  $n$ ; it has the largest estimated mean among all populations that have been sampled at least  $\delta n$  times. The rule  $\varphi^*$ , therefore, compares at stage  $n + 1 = km + j$  the population  $\Pi_j$  with the leader  $\Pi_{j_n}$ . It samples from  $\Pi_j$  if the upper confidence bound  $U_n(j)$  for the mean of  $\Pi_j$  does not fall below the estimated mean of  $\Pi_{j_n}$ ; otherwise it samples from  $\Pi_{j_n}$ . We now establish the asymptotic efficiency of this rule.

**THEOREM 3.** *Assume that  $I(\theta, \lambda)$  satisfies (1.6) and (1.7) and that the functions  $g_{ni}$  and  $h_i$  satisfy (3.1)–(3.5). For  $j = 1, \dots, k$ , let  $T_n(j)$  be the number of times that the rule  $\varphi^*$  samples from  $\Pi_j$  up to stage  $n$ , as defined in (1.2). Define  $\theta^*$  as in (1.4).*

(i) *For every  $\theta = (\theta_1, \dots, \theta_k)$  and every  $j$  such that  $\mu(\theta_j) < \mu(\theta^*)$ ,*

$$E_\theta T_n(j) \leq \left( \frac{1}{I(\theta_j, \theta^*)} + o(1) \right) \log n. \quad (3.9)$$

(ii) *Assume also that  $\Theta$  satisfies (1.9). Then  $E_\theta T_n(j) \sim (\log n)/I(\theta_j, \theta^*)$  for every  $j$  such that  $\mu(\theta_j) < \mu(\theta^*)$ , and the regret of  $\varphi^*$  satisfies (1.8).*

*Proof.* To prove (3.9), let  $L = \{1 \leq l \leq k : \mu(\theta_l) = \mu(\theta^*)\}$ . Let  $0 < \epsilon < \mu(\theta^*) - \max_{j \notin L} \mu(\theta_j)$ . Using the notation  $\#A$  to denote the number of elements of a set  $A$ , we note that for any fixed  $j \notin L$ ,

$$\begin{aligned} T_N(j) \leq & \#\{1 \leq n \leq N - 1 : j_n \in L, |\hat{\mu}_n(j_n) - \mu(\theta^*)| \leq \epsilon, \text{ and} \\ & \varphi^* \text{ samples from } \Pi_j \text{ at stage } n + 1\} + 1 \\ & + \#\{1 \leq n \leq N - 1 : j_n \in L \text{ and } |\hat{\mu}_n(j_n) - \mu(\theta^*)| > \epsilon\} \\ & + \#\{1 \leq n \leq N - 1 : j_n \notin L\}. \end{aligned} \quad (3.10)$$

Let  $Y_{j1}, Y_{j2}, \dots$  denote successive i.i.d. observations from  $\Pi_j$ . From the

definition of  $\varphi^*$  it follows that

$$\begin{aligned}
& \# \{ 2 \leq m \leq N : j_{m-1} \in L, |\hat{\mu}_{m-1}(j_{m-1}) - \mu(\theta^*)| \leq \epsilon, \text{ and} \\
& \quad \varphi^* \text{ samples from } \Pi_j \text{ at stage } m \} \\
& \leq \# \{ 1 \leq \nu \leq N : \varphi^* \text{ samples } Y_{j_\nu} \text{ at some stage } m \text{ with} \\
& \quad \nu \leq m \leq N \text{ and } j_{m-1} \in L, |\hat{\mu}_{m-1}(j_{m-1}) - \mu(\theta^*)| \leq \epsilon \} \\
& \leq 1 + \# \{ 1 \leq i \leq N-1 : g_{Ni}(Y_{j_1}, \dots, Y_{j_i}) \geq \mu(\theta^*) - \epsilon \\
& \quad \text{for some } i \leq n \leq N-1 \} \\
& \leq 1 + \# \{ 1 \leq i \leq N-1 : g_{Ni}(Y_{j_1}, \dots, Y_{j_i}) \geq \mu(\theta^*) - \epsilon \}, \\
& \hspace{25em} \text{by (3.3)}. \quad (3.11)
\end{aligned}$$

In view of (3.2), for every  $\rho > 0$ , we can choose  $\epsilon > 0$  so small that

$$\begin{aligned}
& E_\theta \{ \# \{ 1 \leq i \leq N-1 : g_{Ni}(Y_{j_1}, \dots, Y_{j_i}) \geq \mu(\theta^*) - \epsilon \} \} \\
& \leq \sum_{i=1}^N P_\theta \{ g_{Ni}(Y_{j_1}, \dots, Y_{j_i}) \geq \mu(\theta^*) - \epsilon \} \\
& \leq \frac{1 + \rho + o(1)}{I(\theta_j, \theta^*)} \log N. \quad (3.12)
\end{aligned}$$

Since  $T_n(j_n) \geq \delta n$  by (3.7), it follows that

$$\begin{aligned}
& P_\theta \{ j_n \in L \text{ and } |\hat{\mu}_n(j_n) - \mu(\theta^*)| > \epsilon \} \\
& \leq P_\theta \left\{ \max_{l \in L} \max_{\delta n \leq i \leq n} |h_i(Y_{l_1}, \dots, Y_{l_i}) - \mu(\theta^*)| > \epsilon \right\} \\
& = o(n^{-1}) \quad \text{by (3.5),}
\end{aligned}$$

and therefore

$$E_\theta \{ \# \{ 1 \leq n \leq N-1 : j_n \in L \text{ and } |\hat{\mu}_n(j_n) - \mu(\theta^*)| > \epsilon \} \} = o(\log N). \quad (3.13)$$

It will be shown in Lemma 1 below that

$$E_\theta \{ \# \{ 1 \leq n \leq N-1 : j_n \notin L \} \} = o(\log N). \quad (3.14)$$

From (3.10)–(3.14), (3.9) follows.

If  $\Theta$  also satisfies (1.9), then by Theorem 2,  $E_\theta T_n(j) \geq (1/I(\theta_j, \theta^*) + o(1)) \log n$ . This and (3.9) imply that

$$E_\theta T_n(j) \sim (\log n) / I(\theta_j, \theta^*)$$

for every  $j$  such that  $\mu(\theta_j) < \mu(\theta^*)$ , and (1.8) follows.

LEMMA 1. *With the same notation and assumptions as in Theorem 3(i), let  $L = \{1 \leq l \leq k: \mu(\theta_l) = \mu(\theta^*)\}$ . Let  $0 < \epsilon < \{\mu(\theta^*) - \max_{j \notin L} \mu(\theta_j)\}/2$ , and let  $c$  be a positive integer. For  $r = 0, 1, \dots$ , define*

$$A_r = \bigcap_{1 \leq j \leq k} \left\{ \max_{\delta c^{r-1} \leq n \leq c^{r+1}} |h_n(Y_{j1}, \dots, Y_{jn}) - \mu(\theta_j)| \leq \epsilon \right\},$$

$$B_r = \bigcap_{l \in L} \left\{ g_{ni}(Y_{l1}, \dots, Y_{li}) \geq \mu(\theta^*) - \epsilon \text{ for all } 1 \leq i \leq \delta n \right. \\ \left. \text{and } c^{r-1} \leq n \leq c^{r+1} \right\},$$

where  $0 < \delta < 1/k$  is the same as that used in the rule  $\varphi^*$ . Then

$$(i) P_\theta(\bar{A}_r) = o(c^{-r}), \quad P_\theta(\bar{B}_r) = o(c^{-r}),$$

where  $\bar{A}$  denotes the complement of an event  $A$ . Moreover, if  $c > (1 - k\delta)^{-1}$  and  $r \geq r_0$  (sufficiently large), then

$$(ii) \text{ on } A_r \cap B_r, j_n \in L \text{ for all } c^r \leq n \leq c^{r+1}.$$

Consequently,

$$(iii) E_\theta(\#\{1 \leq n \leq N: j_n \notin L\}) = \sum_{n=1}^N P_\theta\{j_n \notin L\} = o(\log N).$$

*Proof.* (i) From (3.5), it follows that  $P(\bar{A}_r) = o(c^{-r})$ . Let  $[x]$  denote the largest integer  $\leq x$ , and let  $p$  be the smallest positive integer such that  $[c^{r-1}/\delta^p] \geq c^{r+1}$ . For  $t = 0, \dots, p$ , let  $n_t = [c^{r-1}/\delta^t]$ , and define

$$D_t = \bigcap_{l \in L} \left\{ g_{n_t, i}(Y_{l1}, \dots, Y_{li}) \geq \mu(\theta^*) - \epsilon \text{ for all } i \leq n_t \right\}.$$

Then by (3.1),

$$P_\theta(\bar{D}_t) = o(n_t^{-1}) = o(c^{-r}) \quad \text{for } t = 0, \dots, p. \quad (3.15)$$

Given  $c^{r-1} \leq n < c^{r+1}$  and  $1 \leq i \leq \delta n$ , there exists  $t \in \{0, \dots, p-1\}$  such that  $n_{t+1} > n \geq n_t \geq i$ , and therefore by (3.3)

$$g_{ni}(Y_{l1}, \dots, Y_{li}) \geq g_{n_t, i}(Y_{l1}, \dots, Y_{li}) \geq \mu(\theta^*) - \epsilon$$

for all  $l \in L$  on the event  $\bigcap_{0 \leq t \leq p} D_t$ . It then follows that  $B_r \supset \bigcap_{0 \leq t \leq p} D_t$ , and therefore by (3.15),  $P_\theta(\bar{B}_r) = o(c^{-r})$ .

(ii) We now assume that  $(1 - c^{-1})/k > \delta$ . We shall say that at stage  $n$  the rule  $\varphi^*$  samples from  $L$  if it samples from  $\Pi_l$  for some  $l \in L$ . Let

$$v_L(n) = \sum_{l \in L} T_n(l)$$

be the number of times that  $\varphi^*$  samples from  $L$  up to stage  $n$ . We note that

$$\max_{l \in L} T_n(l) \geq v_L(n)/\#L. \quad (3.16)$$

Consider the stage  $n + 1 = km + l$  with  $l \in L$  and  $c^{r-1} \leq n < c^{r+1}$ . We now show that at this stage  $\varphi^*$  must sample from  $L$  on the event  $A_r \cap B_r$ . First note that if  $j_n \in L$ , then  $\varphi^*$  samples from either  $\Pi_{j_n}$  or  $\Pi_l$  at stage  $n + 1 = km + l$ . Now assume that  $j_n \notin L$ . Then since  $T_n(j_n) \geq \delta n$ ,

$$\hat{\mu}_n(j_n) \leq \max_{j \notin L} \mu(\theta_j) + \epsilon < \mu(\theta^*) - \epsilon \quad \text{on } A_r. \quad (3.17)$$

In the case  $T_n(l) \geq \delta n$ , we have on  $A_r$ ,

$$\mu(\theta^*) - \epsilon \leq h_{T_n(l)}(Y_{l_1}, \dots, Y_{l, T_n(l)}) \leq g_{n, T_n(l)}(Y_{l_1}, \dots, Y_{l, T_n(l)}) \quad (3.18)$$

by (3.4), and therefore by (3.17) and (3.18),  $\varphi^*$  samples from  $\Pi_l$  at stage  $n + 1$ . In the case  $T_n(l) < \delta n$ , we have on the event  $B_r$ ,

$$\mu(\theta^*) - \epsilon \leq g_{n, T_n(l)}(Y_{l_1}, \dots, Y_{l, T_n(l)}), \quad (3.19)$$

and therefore by (3.17) and (3.19),  $\varphi^*$  also samples from  $\Pi_l$  at stage  $n + 1$  on  $A_r \cap B_r$ .

On the event  $A_r \cap B_r$ , since  $\varphi^*$  must sample from  $L$  at stage  $n + 1 = km + l$  with  $l \in L$  and  $c^{r-1} \leq n \leq c^{r+1}$ , and since  $(1 - c^{-1})/k > \delta$ , it follows that

$$v_L(n) \geq (\#L/k)(n - c^{r-1} - 2k) > (\#L)\delta n \quad (3.20)$$

for all  $c^r \leq n \leq c^{r+1}$  and  $r \geq r_0$  (sufficiently large). From (3.16) and (3.20), we obtain that on  $A_r \cap B_r$ ,

$$\max_{l \in L} T_n(l) > \delta n \quad \text{for all } c^r \leq n \leq c^{r+1}, \quad (3.21)$$

if  $r \geq r_0$ . We note that for  $r \geq r_0$  and  $c^r \leq n \leq c^{r+1}$ , on the event  $A_r \cap B_r$ ,

$$\begin{aligned} & \max\{\hat{\mu}_n(j) : T_n(j) \geq \delta n \text{ and } j \notin L\} \\ & \leq \max_{j \notin L} \mu(\theta_j) + \epsilon < \mu(\theta^*) - \epsilon \\ & \leq \min\{\hat{\mu}_n(l) : T_n(l) \geq \delta n \text{ and } l \in L\}, \end{aligned}$$

the last set being nonempty by (3.21). Hence  $j_n \in L$  for all  $c^r \leq n \leq c^{r+1}$  on  $A_r \cap B_r$  if  $r \geq r_0$ .

(iii) Let  $c > (1 - k\delta)^{-1}$ . Then it follows from (i) and (ii) that for  $r \geq r_0$  and  $c^r \leq n \leq c^{r+1}$ ,

$$P_\theta\{j_n \notin L\} \leq P_\theta(\bar{A}_r) + P_\theta(\bar{B}_r) = o(c^{-r}),$$

and therefore  $\sum_{c^r \leq n \leq c^{r+1}} P_\theta\{j_n \notin L\} = o(1)$ . Hence  $\sum_{n=1}^N P_\theta\{j_n \notin L\} = o(\log N)$ .

#### 4. CONFIDENCE SEQUENCES AND ALLOCATION RULES FOR SPECIAL DISTRIBUTIONS

In this section we make use of certain generalized likelihood ratios to construct confidence sequences that satisfy the conditions (3.1)–(3.3). These generalized likelihood ratios are described in the following lemma.

LEMMA 2. Let  $Y_1, Y_2, \dots$  be i.i.d. random variables with a common density  $f(y; \theta)$  with respect to some measure  $\nu$ , where  $\theta$  is a real parameter.

(i) Let  $\hat{\theta}_n = \hat{\theta}_n(Y_1, \dots, Y_n)$  be an estimate of  $\theta$  at stage  $n$ . Then for every  $a \geq 1$ ,

$$P_\theta \left\{ \prod_{i=1}^n f(Y_i; \hat{\theta}_{i-1}) / \prod_{i=1}^n f(Y_i; \theta) \geq a \text{ for some } n \geq 1 \right\} \leq a^{-1}. \quad (4.1)$$

(ii) Let  $C$  be a compact set of real numbers such that for every  $\lambda \in C$  and some  $\theta$

$$\lim_{\delta \downarrow 0} E_\theta(\sup\{f(Y_1; r)/f(Y_1; \theta) : r \in C, |r - \lambda| < \delta\}) = 1. \quad (4.2)$$

Then for every  $d > 0$ ,

$$\limsup_{a \rightarrow \infty} (\log a)^{-1} \log P_\theta \left\{ \sup_{\lambda \in C} \prod_{i=1}^n f(Y_i; \lambda) / \prod_{i=1}^n f(Y_i; \theta) \geq a \right. \\ \left. \text{for some } n \leq d \log a \right\} \leq -1. \quad (4.3)$$

Moreover, for every  $d > 0$  and  $0 < \rho < 1$ ,

$$\limsup_{a \rightarrow \infty} (\log a)^{-1} \log P_\gamma \left\{ \prod_{i=1}^n f(Y_i; \gamma) / \prod_{i=1}^n f(Y_i; \theta) \leq \rho^n \right. \\ \left. \text{and } \sup_{\lambda \in C} \prod_{i=1}^n f(Y_i; \lambda) / \prod_{i=1}^n f(Y_i; \theta) \geq a \text{ for some } n \geq d \log a \right\} \\ \leq -1 + d \log \rho < -1. \quad (4.4)$$

*Proof.* (i) Under  $P_\theta$ , the sequence  $\{\prod_{i=1}^n f(Y_i; \hat{\theta}_{i-1}) / \prod_{i=1}^n f(Y_i; \theta), n \geq 1\}$  is a nonnegative martingale with mean 1, and therefore (4.1) follows (cf. [4]).

(ii) To prove (4.3), in view of (4.2) we can choose for every  $\epsilon > 0$  and  $\lambda \in C$  a positive constant  $\delta(\epsilon, \lambda)$  such that

$$E_{\theta} \left( \sup_{r \in B(\lambda)} f(Y_1; r) / f(Y_1; \theta) \right) < 1 + \epsilon, \quad (4.5)$$

where  $B(\lambda) = \{r \in C: |r - \lambda| < \delta(\epsilon, \lambda)\}$ . From (4.5) it follows that

$$P_{\theta} \left\{ \sup_{r \in B(\lambda)} \prod_{i=1}^n f(Y_i; r) / \prod_{i=1}^n f(Y_i; \theta) \geq a \right\} \leq a^{-1}(1 + \epsilon)^n.$$

Since  $C$  is compact, we can choose a finite covering  $B(\lambda_1), \dots, B(\lambda_m)$  of  $C$ , and therefore

$$P_{\theta} \left\{ \sup_{r \in C} \prod_{i=1}^n f(Y_i; r) / \prod_{i=1}^n f(Y_i; \theta) \geq a \right\} \leq ma^{-1}(1 + \epsilon)^n. \quad (4.6)$$

Since  $\epsilon$  can be arbitrarily small, (4.3) follows from (4.6).

To prove (4.4), let  $F_n = \{\prod_{i=1}^n f(Y_i; \gamma) / \prod_{i=1}^n f(Y_i; \theta) \leq \rho^n\}$ ,  $G_n = \{\sup_{\lambda \in C} \prod_{i=1}^n f(Y_i; \lambda) / \prod_{i=1}^n f(Y_i; \theta) \geq a\}$ . We note that

$$\begin{aligned} P_{\gamma}(F_n \cap G_n) &= \int_{F_n \cap G_n} \prod_{i=1}^n (f(Y_i; \gamma) / f(Y_i; \theta)) dP_{\theta} \\ &\leq \rho^n P_{\theta}(G_n) \leq ma^{-1} \{\rho(1 + \epsilon)\}^n, \quad \text{by (4.6)}. \end{aligned} \quad (4.7)$$

Choosing  $\epsilon$  so small that  $\rho(1 + \epsilon) < 1$ , we have

$$\sum_{n \geq d \log a} \{\rho(1 + \epsilon)\}^n = O(\{\rho(1 + \epsilon)\}^{d \log a}),$$

and therefore (4.4) follows from (4.7).

We now apply the preceding results to construct confidence sequences and asymptotically efficient allocation rules for normal, Bernoulli, Poisson, and double exponential populations.

**EXAMPLE 1.** Let  $Y_{ji}$ ,  $j = 1, 2, \dots, k$ ,  $i = 1, 2, \dots$ , be independent normal random variables with known common variance  $\sigma^2 > 0$  and unknown means  $EY_{ji} = \theta_j$ . Thus,  $\mu(\theta) = \theta$ ,  $\Theta = (-\infty, \infty)$ ,  $\nu =$  Lebesgue measure,

$$f(y, \theta) = (2\pi\sigma^2)^{-1/2} \exp\{- (y - \theta)^2 / 2\sigma^2\}, \quad (4.8)$$

and

$$I(\theta, \lambda) = (\theta - \lambda)^2 / (2\sigma^2). \quad (4.9)$$

Conditions (1.6), (1.7), and (1.9) are clearly satisfied.

Let  $a_{ni}$  ( $n = 1, 2, \dots, i = 1, \dots, n$ ) be positive constants such that for every fixed  $i$

$$a_{ni} \text{ is nondecreasing in } n \geq i, \quad (4.10)$$

and there exist  $\epsilon_n \rightarrow 0$  for which

$$|a_{ni} - (\log n)/i| \leq \epsilon_n (\log n)^{1/2}/i^{1/2} \text{ for all } i \leq n. \quad (4.11)$$

For  $j = 1, \dots, k$ , define

$$\bar{Y}_i(j) = (Y_{j1} + \dots + Y_{ji})/i, \quad h_i(Y_{j1}, \dots, Y_{ji}) = \bar{Y}_i(j), \quad (4.12)$$

$$g_{ni}(Y_{j1}, \dots, Y_{ji}) = \bar{Y}_i(j) + \sigma(2a_{ni})^{1/2} \quad \text{for } n \geq i. \quad (4.13)$$

Obviously, for every  $0 < \delta < 1$  and  $\epsilon > 0$ ,

$$P\left\{ \max_{\delta n \leq i \leq n} |\bar{Y}_i(j) - \theta_j| > \epsilon \right\} = o(n^{-1}), \quad (4.14)$$

so condition (3.5) is satisfied. From (4.11) and the tail probability of the normal distribution it follows easily that for  $r < \theta_j$

$$\sum_{i=1}^n P\{r > \bar{Y}_i(j) + \sigma(2a_{ni})^{1/2}\} = o(n^{-1}),$$

and therefore condition (3.1) is satisfied.

Conditions (3.3) and (3.4) are obviously satisfied. We now show that (3.2) also holds. Let  $\lambda > \theta_j$  and define

$$L_n = \sup\{1 \leq i \leq n: \bar{Y}_i(j) + \sigma(2a_{ni})^{1/2} \geq \lambda\} \quad (\sup \emptyset = 0),$$

$$T_\epsilon = \sup\{i \geq 1: |\bar{Y}_i(j) - \theta_j| \geq \epsilon\}.$$

Then  $ET_\epsilon < \infty$  for all  $\epsilon > 0$  (cf. [1]). Moreover, it follows from (4.11) that for  $0 < \epsilon < \lambda - \theta_j$

$$E(L_n I_{\{L_n > T_\epsilon\}}) \leq 2\sigma^2(\lambda - \theta_j - \epsilon)^{-2}(1 + o(1))\log n \quad \text{as } n \rightarrow \infty.$$

Obviously,  $E(L_n I_{\{L_n \leq T_\epsilon\}}) \leq ET_\epsilon$ . By the strong law of large numbers and Fatou's lemma,

$$EL_n \geq 2\sigma^2(\lambda - \theta_j)^{-2}(1 + o(1))\log n.$$

Hence, letting  $\epsilon \downarrow 0$ , we obtain that

$$EL_n \sim 2\sigma^2(\lambda - \theta_j)^{-2}\log n = (\log n)/I(\theta_j, \lambda). \quad (4.15)$$

Since  $\#\{1 \leq i \leq n: \bar{Y}_i(j) + \sigma(2a_{ni})^{1/2} \geq \lambda\} \leq L_n$ , it follows from (4.15)

that for  $\lambda > \theta_j$

$$\sum_{i=1}^n P\{\bar{Y}_i(j) + \sigma(2a_{ni})^{1/2} \geq \lambda\} \leq (1 + o(1))(\log n)/I(\theta_j, \lambda).$$

With  $h_i$  and  $g_{ni}$  given by (4.12) and (4.13), we define the allocation rule  $\varphi^*$  as in Section 3. Theorem 3 is therefore applicable to this special case and shows that  $\varphi^*$  provides an asymptotically efficient allocation rule for  $k$  normal populations with common known variance  $\sigma^2$ . We note that  $\bar{Y}_i(j)$  is the maximum likelihood estimate of  $\theta_j$  based on  $Y_{j1}, \dots, Y_{ji}$ , and that the upper confidence bound  $g_{ni}(Y_{j1}, \dots, Y_{j1})$  can be expressed in terms of generalized likelihood ratios as follows:

$$\begin{aligned} g_{ni}(Y_{j1}, \dots, Y_{ji}) &= \inf\{\lambda \geq \bar{Y}_i(j) : I(\bar{Y}_i(j), \lambda) \geq a_{ni}\} \\ &= \inf\left\{\lambda \geq \bar{Y}_i(j) : \sup_{\theta} \prod_{i=1}^i f(Y_{ji}; \theta) / \prod_{i=1}^i f(Y_{ji}; \lambda) \geq e^{ia_{ni}}\right\}. \end{aligned} \quad (4.16)$$

The upper confidence bounds in the next two examples are also of this general form.

**EXAMPLE 2.** Let  $Y_{ji}$  be independent Bernoulli random variables such that  $Y_{ji}$  has density

$$f(y; \theta_j) = \theta_j^y (1 - \theta_j)^{1-y}, \quad y = 0, 1, \quad (4.17)$$

with respect to the counting measure  $\nu$ . Here  $\mu(\theta) = \theta$ ,  $\Theta = (0, 1)$ , and

$$I(\theta, \lambda) = \theta \log(\theta/\lambda) + (1 - \theta) \log\{(1 - \theta)/(1 - \lambda)\}. \quad (4.18)$$

Conditions (1.6), (1.7), and (1.9) are clearly satisfied.

For  $j = 1, \dots, k$ , define  $\bar{Y}_i(j)$  and  $h_i$  as in (4.12), and note that (4.14) still holds. Let  $a_{ni}$  ( $n = 1, 2, \dots$ ,  $i = 1, \dots, n$ ) be positive constants satisfying (4.10) and such that

$$\lim_{\delta \downarrow 0} \left( \liminf_{n \rightarrow \infty} \min\{a_{ni} : i \leq \delta \log n\} \right) = \infty, \quad (4.19a)$$

$$\lim_{d \rightarrow \infty} \left( \limsup_{n \rightarrow \infty} \max\{a_{ni} : d \log n \leq i \leq n\} \right) = 0, \quad (4.19b)$$

and

$$\lim_{n \rightarrow \infty} \max\{|ia_{ni}/\log n - 1| : \delta \log n \leq i \leq d \log n\} = 0 \quad \forall 0 < \delta < d. \quad (4.19c)$$

(Note that (4.19) is a weaker assumption than (4.11).) Define  $g_{ni}$  as in

(4.16), where the parameter  $\lambda$  can only take values in  $(0, 1)$  and where we now set  $\inf \emptyset = 1$ . Since  $I(\bar{Y}_i(j), \lambda)$  is a convex function in  $\lambda$  with minimum at  $\lambda = \bar{Y}_i(j)$ , we have the equivalence

$$1 > r \geq g_{ni}(Y_{j1}, \dots, Y_{ji}) \Leftrightarrow 1 > r \geq \bar{Y}_i(j) \text{ and } I(\bar{Y}_i(j), r) \geq a_{ni}. \quad (4.20)$$

Let  $1 > \lambda > \theta_j$  and define  $L = \sup\{i: \bar{Y}_i(j) \geq \lambda\}$ ,  $L_n = \sup\{1 \leq i \leq n: I(\bar{Y}_i(j), \lambda) < a_{ni}\}$ . Then  $EL < \infty$  (cf. [1]), and an argument similar to the proof of (4.15) shows that  $EL_n \sim (\log n)/I(\theta_j, \lambda)$ . From (4.20), it follows that  $\#\{1 \leq i \leq n: g_{ni}(Y_{j1}, \dots, Y_{ji}) > \lambda\} \leq L + L_n$ , and therefore

$$\sum_{i=1}^n P\{g_{ni}(Y_{j1}, \dots, Y_{ji}) > \lambda\} \leq (1 + o(1))(\log n)/I(\theta_j, \lambda).$$

Hence condition (3.2) is satisfied.

We now show that condition (3.1) is also satisfied. Let  $\theta_j > r > 0$ . Then by (4.19a), we can choose  $\delta > 0$  and  $n_0$  such that

$$\sup_{\theta} I(\theta, r) = \max\{|\log r|, |\log(1 - r)|\} < a_{ni} \quad \text{for } i \leq \delta \log n \text{ and } n \geq n_0. \quad (4.21)$$

Let  $\rho = (\theta_j/r)^r \{(1 - \theta_j)/(1 - r)\}^{1-r}$ . We note that  $\log \rho = -I(r, \theta_j) < 0$  and that

$$r \geq \bar{Y}_i(j) \Rightarrow \prod_{i=1}^i f(Y_{ji}; \theta_j) / \prod_{i=1}^i f(Y_{ji}; r) \leq \rho^i. \quad (4.22)$$

Hence  $P\{r \geq \bar{Y}_i(j)\} \leq \rho^i$ , so we can choose  $d > \delta$  such that

$$P\{r \geq \bar{Y}_i(j) \text{ for some } i \geq d \log n\} = o(n^{-1}).$$

Moreover, from (4.20), (4.21), and (4.22), it follows that

$$\begin{aligned} & P\{r \geq g_{ni}(Y_{j1}, \dots, Y_{ji}) \text{ for some } i \leq d \log n\} \\ &= P\{r \geq \bar{Y}_i(j) \text{ and } I(\bar{Y}_i(j), r) \geq a_{ni} \text{ for some } \delta \log n \leq i \leq d \log n\} \\ &\leq P\left\{ \prod_{i=1}^i f(Y_{ji}; \theta_j) / \prod_{i=1}^i f(Y_{ji}; r) \leq \rho^i \text{ and } \right. \\ &\quad \left. \sup_{0 \leq \theta \leq r} \prod_{i=1}^i f(Y_{ji}; \theta) / \prod_{i=1}^i f(Y_{ji}; r) \geq e^{ia_{ni}} \right. \\ &\quad \left. \text{for some } \delta \log n \leq i \leq d \log n \right\} \\ &= o(n^{-1}), \quad \text{by (4.19c) and Lemma 2 (ii).} \end{aligned}$$

With  $h_i$  defined in (4.12) and  $g_{ni}$  defined in (4.16), the argument above shows that Theorem 3 is applicable, so the rule  $\varphi^*$  provides an asymptotically efficient allocation rule for  $k$  Bernoulli populations. In view of (4.20), we do not need the explicit value of  $g_{ni}(Y_{j1}, \dots, Y_{ji})$  to implement the rule  $\varphi^*$ . In fact, the allocation criterion (3.8) can now be rewritten as follows: Letting  $\hat{\mu}_n(j) = \bar{Y}_{T_n(j)}(j)$ , sample from  $\Pi_j$  at stage  $n + 1 = km + j$  only if

$$\hat{\mu}_n(j) \geq \hat{\mu}_n(j_n) \quad \text{or} \quad I(\hat{\mu}_n(j), \hat{\mu}_n(j_n)) \leq a_{n, T_n(j)}. \quad (4.23)$$

**EXAMPLE 3.** Let  $Y_{ji}$  be independent Poisson random variables such that  $Y_{ji}$  has density

$$f(y; \theta_j) = e^{-\theta_j} \theta_j^y / y!, \quad y = 0, 1, \dots$$

with respect to the counting measure  $\nu$ . Here  $\mu(\theta) = \theta$ ,  $\Theta = (0, \infty)$ , and

$$I(\theta, \lambda) = \theta \log(\theta/\lambda) - (\theta - \lambda).$$

Conditions (1.6), (1.7), and (1.9) are clearly satisfied. Letting  $a_{ni}$  be positive constants satisfying (4.10) and (4.19), and defining  $h_i$  as in (4.12) and  $g_{ni}$  as in (4.16), we can use the argument of Example 2 to show that Theorem 3 is again applicable, noting that  $\sup_{0 \leq \theta \leq r} I(\theta, r) = r$ . Moreover, the allocation criterion (3.8) of the rule  $\varphi^*$  of Theorem 3 can be written in the more convenient form (4.23).

In each of the preceding examples the sample means of the  $k$  populations are sufficient statistics and are used in the allocation rule  $\varphi^*$ . We now give an example in which no simple sufficient statistics are available and in which sample medians are used instead of sample means. Moreover, in this example we show how the generalized likelihood ratios of Lemma 2(i) can be applied to construct confidence sequences satisfying (3.1).

**EXAMPLE 4.** Let  $Y_{ji}$  be independent double exponential random variables such that  $Y_{ji}$  has density

$$f(y; \theta_j) = \frac{1}{2} \exp(-|y - \theta_j|), \quad -\infty < y < \infty. \quad (4.24)$$

Here  $\mu(\theta) = \theta$ ,  $\Theta = (-\infty, \infty)$ ,  $\nu$  is Lebesgue measure, and  $I(\theta, \lambda) = |\theta - \lambda|$ . Conditions (1.6), (1.7), and (1.9) are clearly satisfied.

Let  $b_n$  be a nondecreasing sequence of positive numbers such that

$$b_n \rightarrow \infty \quad \text{and} \quad \log b_n = o(\log n) \quad \text{as } n \rightarrow \infty. \quad (4.25)$$

For  $j = 1, \dots, k$  define

$$M_i(j) = \text{med}\{Y_{j1}, \dots, Y_{ji}\}, \quad h_i(Y_{j1}, \dots, Y_{ji}) = M_i(j), \quad (4.26)$$

$$g_{ni}(Y_{j1}, \dots, Y_{ji}) = \inf \left\{ \theta \geq M_i(j) : \sum_{t=1}^i |Y_{jt} - \theta| \geq \log nb_n + \sum_{t=1}^i |Y_{jt} - M_{t-1}(j)| \right\}. \quad (4.27)$$

Since  $u(\theta) = \sum_{t=1}^i |Y_{jt} - \theta|$  is a piecewise linear and convex function of  $\theta$  with its minimum at  $\theta = M_i(j)$ , we have the equivalence

$$\begin{aligned} \theta &\geq g_{ni}(Y_{j1}, \dots, Y_{ji}) \\ \Leftrightarrow \theta &\geq M_i(j) \text{ and } \sum_{t=1}^i |Y_{jt} - \theta| \geq \log nb_n + \sum_{t=1}^i |Y_{jt} - M_{t-1}(j)|. \end{aligned} \quad (4.28)$$

Since for every  $\epsilon > 0$ ,

$$P\{|M_i(j) - \theta_j| > \epsilon\} = O(\rho^i) \quad \text{for some } 0 < \rho = \rho(\epsilon) < 1, \quad (4.29)$$

it follows that condition (3.5) is satisfied. Moreover, conditions (3.3) and (3.4) are also satisfied. Furthermore, for  $r < \theta_j$ ,

$$\begin{aligned} &P\{r \geq g_{ni}(Y_{j1}, \dots, Y_{ji}) \text{ for some } i \leq n\} \\ &\leq P\{\theta_j \geq g_{ni}(Y_{j1}, \dots, Y_{ji}) \text{ for some } i \leq n\} \\ &\leq P\left\{ \prod_{t=1}^i f(Y_{jt}; M_{t-1}(j)) / \prod_{t=1}^i f(Y_{jt}; \theta_j) \geq nb_n \text{ for some } i \geq 1 \right\}, \\ &\quad \text{by (4.24) and (4.28),} \\ &\leq (nb_n)^{-1}, \quad \text{by Lemma 2(i),} \\ &= o(n^{-1}). \end{aligned}$$

Hence condition (3.1) is satisfied.

Using the results of [1], it can be shown that condition (3.2) is also satisfied. Hence, with  $h_i$  and  $g_{ni}$  given by (4.26) and (4.27), Theorem 3 can be applied to show that  $\varphi^*$  is an asymptotically efficient allocation rule for

double exponential populations. In view of (4.28), the allocation criterion (3.8) can be rewritten as follows: Sample from  $\Pi_j$  at stage  $n + 1 = km + j$  only if

$$M_{T_n(j)}(j) \geq M_{T_n(j_n)}(j_n) \quad (4.30)$$

or

$$\sum_{i=1}^{T_n(j)} |Y_{ji} - M_{T_n(j_n)}(j_n)| \leq \log nb_n + \sum_{i=1}^{T_n(j)} |Y_{ji} - M_{i-1}(j)|.$$

#### REFERENCES

1. Y. S. CHOW AND T. L. LAI, Some one-sided theorems on the tail distribution of sample sums with applications to the last time and largest excess of boundary crossings, *Trans. Amer. Math. Soc.* **208** (1975), 51-72.
2. H. REIMNITZ, Asymptotic near admissibility and asymptotic near optimality in the two armed bandit problem, *Math. Operationsforsch. Statist. Ser. Statist.*, in press.
3. H. ROBBINS, Some aspects of the sequential design of experiments, *Bull. Amer. Math. Soc.* **55** (1952), 527-535.
4. H. ROBBINS AND D. SIEGMUND, A class of stopping rules for testing parametric hypotheses, *Proc. Sixth Berkeley Symp. Math. Statist. Probab.* **4** (1972), 37-41.